

ReQoL

Recovering Quality of Life



Anju Keetharuth, Janice Connell, Milad Karimi, Michael Barkham, Tom Ricketts and John Brazier, University of Sheffield

What is Recovering Quality of Life (ReQoL)?

- ✓ NEW generic Patient Reported Outcome Measure to assess Quality of Life in people with mental health conditions
- ✓ SHORT service user outcome measure for use in **routine practice**
- ✓ Developed from **service users' perspectives** with inputs from clinicians, national and international experts
- ✓ Commissioned and funded by the Department of Health


Why ReQoL?

- ✓ Existing generic measures (EQ-5D and SF-36) are not suitable for use with mental health service users
- ✓ An appropriate outcome measure is necessary for use in the NHS Outcomes Framework
- ✓ An appropriate outcome measure is recommended in the context of the national tariff payment system (PbR)

Stage 1: Identifying Domains **Method:** The domains of the questionnaire were identified through a systematic review of the quality of life literature and 19 qualitative interviews with service users.

Stage 2: Generating item pool **Method:** A systematic approach was adopted to generate an item pool. The criteria from Norman & Streiner (1989) were used to select items.

Results of stage 1: Seven domains are important for quality of life to people with mental health problems. The domains are similar to factors which are important for Personal Recovery – Connectedness, Hope, Identity, Meaning, and Empowerment (CHIME -Leamy et al., 2011).



Stage 2 Selection Flowchart:


Initial round of selection

N = 1597

ReQoL Sheffield team

N = 122

N = 180



- Expert User Group 123
 - 122 items were presented
 Expert users added 58 items yielding 180 items

item was recorded

Scientific Group N = 101

ReQoL Sheffield team
N = 87

- Reviewed all 180 items
 Applied Norman &
 Streiner's criteria (reading age, jargon, value-laden, length of items, double negatives ...)
- Ensured items covered all sub themes and severity levels

Stage 3: Face and content validity

- The items from Stage 2 were tested for face and content validity through 59 interviews with service users.
- Additional inputs on the items were obtained from clinicians, experts from our Stakeholder and Advisory Groups, and a translatability assessment.
- Number of items was reduced to 60, some which were rephrased based on comments received.

What next?

PSYCHOMETRICS ANALYSIS: Between March and May 2015, the ReQoL questionnaire will be administered by at least 11 NHS trusts, 6 GP practices, and 4 charities nationally. The data collected will be analysed (factor analysis and Rasch analysis) to produce the final questionnaires: the full ReQoL (30 items) and the short ReQoL (10 items). Both versions will be available for use by August 2015.

Further work

The measure will then be validated (Sept 15 – Feb 2016) and preference weights will also be produced so that the ReQoL questionnaire can be used in economic evaluations (March – May 2016).